

Fragmented many-body ground states for scalar bosons in a single trap

Philipp Bader and Uwe R. Fischer

*Eberhard-Karls-Universität Tübingen, Institut für Theoretische Physik
Auf der Morgenstelle 14, D-72076 Tübingen, Germany*

We investigate whether the many-body ground states of bosons in a generalized two-mode model with localized inhomogeneous single-particle orbitals and anisotropic long-range interactions (e.g. dipole-dipole interactions), are coherent or fragmented. It is demonstrated that fragmentation can take place in a single trap for positive values of the interaction couplings, implying that the system is potentially stable. Furthermore, the degree of fragmentation is shown to be insensitive to small perturbations on the single-particle level.

Fragmentation in many-body states of interacting bosons is defined by the single-particle density matrix having more than one macroscopic eigenvalue [1, 2], leading to macroscopic occupation of more than one state. In the continuum, and for repulsive contact interactions, the Fock term in the total energy favors a single condensate; a negative contact interaction implies the collapse of the gas before ground-state fragmentation sets in [3]. On the other hand, in spatially well-separated systems, fragmentation can be obtained for deep double wells [4], or in its periodic extension Mott state in optical lattices [5]. Further examples for fragmentation have been found [6], e.g., when internal degrees of freedom and sufficient symmetry in the interaction are present [7], in the Richardson pairing model [8], in the crossover to fermionization and for excited states [9], and in rapidly rotating gases [10, 11]. However, no ground-state fragmentation of a scalar Bose gas in a single trap has been reported.

So far, only contact interactions between the bosons and quasi-homogeneous or spatially periodic systems were considered. The question then arises whether for general inhomogeneity, nonlocality and anisotropy, i.e. given the localization in a single trap with inhomogeneously distributed single-particle orbitals (modes) and for partly positive and negative long-range interactions (of which the archetype is the dipole-dipole interaction), ground-state fragmentation can occur. In the following, we consider for simplicity and easy comparison with the double-well case, that only two single-particle orbitals are occupied [12]. Examples are gases strongly confined in a given direction, e.g., close to the quasi-2D limit; the question then arises whether fragmentation occurs with respect to that strongly confining direction. A two-mode approximation has, furthermore, the important benefit that the many-body states of the system can be found essentially exactly numerically (and for special cases analytically). We demonstrate that by tuning the four interaction matrix elements relative to each other, various many-body states – coherent and fragmented states, as well as coherent superpositions of degenerate macroscopically distinct quantum states [13] – can be accessed in a single trap. In addition, the fragmented states are not susceptible to decay to a non-fragmented (coherent) state

because of a perturbation on single-particle level [16, 17], due to the fragmentation being based to the values of the interaction couplings.

We begin with a general quadratic plus quartic Hamiltonian for two interacting modes,

$$\begin{aligned} \hat{H} = & \epsilon_0 \hat{a}_0^\dagger \hat{a}_0 + \epsilon_1 \hat{a}_1^\dagger \hat{a}_1 - \frac{\Omega}{2} (\hat{a}_0^\dagger \hat{a}_1 + \text{h.c.}) \\ & + \frac{A_1}{2} \hat{a}_0^\dagger \hat{a}_0^\dagger \hat{a}_0 \hat{a}_0 + \frac{A_2}{2} \hat{a}_1^\dagger \hat{a}_1^\dagger \hat{a}_1 \hat{a}_1 \\ & + \frac{A_3}{2} (\hat{a}_0^\dagger \hat{a}_0^\dagger \hat{a}_1 \hat{a}_1 + \text{h.c.}) + \frac{A_4}{2} \hat{a}_1^\dagger \hat{a}_1 \hat{a}_0^\dagger \hat{a}_0. \end{aligned} \quad (1)$$

The interaction coefficients are given by $A_1 = V_{0000}, A_2 = V_{1111}, A_3 = V_{1100} = V_{0011}$ (taken to be real), and $A_4 = V_{0101} + V_{1010} + V_{1001} + V_{0110}$, where $V_{ijkl} = \int d^3r \int d^3r' \Psi_i^*(\mathbf{r}) \Psi_j^*(\mathbf{r}') V_{\text{int}}(\mathbf{r} - \mathbf{r}') \Psi_k(\mathbf{r}') \Psi_l(\mathbf{r})$. It is of major importance for the discussion to follow that we include pair-exchange between the two modes due to scattering of pairs of bosons $\propto A_3$, in addition to the standard density-density type terms $\propto A_1, A_2, A_4$. We stress that this term is *absent* in the homogeneous continuum due to momentum conservation. We also note that the independence of the coefficients A_i makes (1) different from rotationally invariant interaction Hamiltonians in gases with internal degrees of freedom, where the modes are components of a spinor [6, 7]. For Fourier transformable interaction potentials we have, by convolution, $V_{ijkl} = \frac{1}{(2\pi)^3} \int d^3k \tilde{\rho}_{il}(-\mathbf{k}) \tilde{V}_{\text{int}}(\mathbf{k}) \tilde{\rho}_{jk}(\mathbf{k})$, where the $\tilde{\rho}_{jk}(\mathbf{k})$ are Fourier transforms of $\Psi_j^*(\mathbf{r}) \Psi_k(\mathbf{r})$. The single-particle energies $\epsilon_i = \int d^3r [\frac{\hbar^2}{2m} |\nabla \Psi_i|^2 + V_{\text{trap}} |\Psi_i|^2]$ are given by kinetic plus trap contributions. Finally, the Josephson-type coupling Ω connects the single-particle states and corresponds to a tunneling rate [18, 19].

We emphasize that differing magnitudes and signs of the A_i can stem both from the interaction *and* the form of the single particle orbitals. This only simplifies in the continuum limit (the orbitals are plane waves then), where the interaction couplings directly reflect anisotropy and long-range character of the interaction. Note that for contact interactions, all A_i negative necessarily implies collapse for large systems, occurring by the fact that choosing ever more singular single particle orbitals is energetically advantageous. On the other hand, by ad-

justing the single-particle orbitals (the trapping) in conjunction with in general anisotropic interactions like the dipole-dipole interaction, the magnitude and sign of the interaction coefficients A_i can be engineered [20]. The Hamiltonian (1) therefore represents a minimal model to investigate whether ground-state fragmentation takes place for potentials and traps with arbitrary anisotropy and shape, leading to essentially independent interaction couplings A_i . In a single trap, the two terms in the last line of (1) are important, in distinction to the double-well case ($\hat{a}_0, \hat{a}_1$ are then the annihilation operators for particles in the left and right wells, respectively), where they both are exponentially suppressed. In particular, we will see that the pair-exchange term $\propto A_3$ decides upon the class of many-body solutions obtained.

Let $|l\rangle \equiv |N-l, l\rangle$ be a N -particle two-mode state, with $N-l$ in the ground and l particles in the excited state. Then, all many-particle states can be expressed by $|\Psi\rangle = \sum_{l=0}^N \psi_l |l\rangle$. The energy reads with the Hamiltonian (1),

$$\langle \Psi | \hat{H} | \Psi \rangle = \sum_{l=0}^N \left[\frac{A_3}{2} d_l (\psi_l^* \psi_{l+2} + \psi_l \psi_{l+2}^*) - \frac{\Omega}{2} \omega_l (\psi_l^* \psi_{l+1} + \psi_l \psi_{l+1}^*) + c_l |\psi_l|^2 \right], \quad (2)$$

where the diagonal coefficient $c_l = \epsilon_0(N-l) + \epsilon_1 l + \frac{1}{2} A_1(N-l)(N-l-1) + \frac{1}{2} A_2 l(l-1) + \frac{1}{2} A_4(N-l)l$, $d_l = \sqrt{(l+2)(l+1)(N-l-1)(N-l)}$, while $\omega_l = \sqrt{(N-l)(l+1)}$. Differentiating the functional (2) with respect to ψ_l , the linear system to be solved reads

$$\langle l | \hat{H} | \Psi \rangle = E \psi_l = \frac{A_3}{2} (d_l \psi_{l+2} + d_{l-2} \psi_{l-2}) - \frac{\Omega}{2} (\omega_l \psi_{l+1} + \omega_{l-1} \psi_{l-1}) + c_l \psi_l. \quad (3)$$

There are three important cases, $A_3 < 0$, $A_3 > 0$, $A_3 = 0$, for a given set of interaction couplings A_1, A_2, A_4 (which enter c_l), assuming $\Omega \geq 0$.

To get first insight, and introduce some notions, we begin by discussing the simplest case of no pair-exchange between orbitals and no coupling of levels, $A_3 = \Omega = 0$. For this configuration, exactly one state $|l\rangle$ will obtain, i.e., a definite occupation of both the ground and excited single-particle state, because the Hamiltonian is diagonal in l . The necessary condition for such a *fragmented* state is convexity of the energy parabola in l , $\frac{\partial^2 c_l}{\partial l^2} = A_1 + A_2 - A_4 > 0$. The energy is minimized for $l_{\min} = \frac{N}{2} + \frac{(A_1 - A_2)(N-1) + 2(\epsilon_0 - \epsilon_1)}{2(A_1 + A_2 - A_4)}$. On the other hand, if the energy parabola is concave, we get a condensate, i.e. all particles accumulating in one of the two-mode states $|N, 0\rangle$ ($l = 0$), $|0, N\rangle$ ($l = N$), depending on which of the two states has lower energy. There is however one exceptional case. If the energies of the two states are identical, leading, by $c_0 = c_N$, to the condition




FIG. 1: Numerical solutions of (4) for the indicated parameter set, for $A_3 = -1$ (left) and $A_3 = 1$ (right), centered around $l = N/2$ (the A_i are given in an arbitrary overall unit of energy). The red squares indicate the even l sector (distribution of the ϕ_l), and the blue circles the odd l sector (distribution of the Φ_l), using the ground-state superposition $|\phi\rangle - |\Phi\rangle$ for clarity of representation.

$\epsilon_1 - \epsilon_0 = (N-1)(A_2 - A_1)/2$, concavity leads to the coherent superposition of the macroscopically distinct and degenerate many-body states $|N, 0\rangle$ and $|0, N\rangle$ [6, 14, 15], characterized by number fluctuations of, say, the lower single-particle state, $\Delta N_0^2 = \langle \hat{N}_0^2 \rangle - \langle \hat{N}_0 \rangle^2 = \mathcal{O}(N^2)$.

In the case of a still vanishing $\Omega = 0$, but with a general set of $\{A_i\}$, Eq. (3) decomposes into two independent equations connecting the even and odd l sectors of ψ_l ,

$$E \psi_{2k} = c_{2k} \psi_{2k} + d_{2k} \psi_{2k+2} + d_{2k-2} \psi_{2k-2}, \\ E \psi_{2k+1} = c_{2k+1} \psi_{2k+1} + d_{2k+1} \psi_{2k+3} + d_{2k-1} \psi_{2k-1}, \quad (4)$$

$\forall k \in \{0, \dots, N/2\}$. We have numerically established that (to faster than exponential accuracy in N , i.e. $\propto \exp[-N^\alpha]$ with $\alpha > 1$) the two resulting ground states in the sectors of even and odd l , $|\phi\rangle = \sum_l \phi_l |2l\rangle$ and $|\Phi\rangle = \sum_l \Phi_l |2l+1\rangle$ are degenerate when the continuum limit (see below) is valid. We display typical results for ψ_l distributions in Fig. 1, where we put $\epsilon_0 = \epsilon_1 = 0$ here and in the plots to follow (the ϵ_i influencing the location of the maximum of the ψ_l distribution, see Eq. (7) and the discussion following it).

When the pair-exchange coupling A_3 is positive (Fig. 1, right panel), it causes alternating signs, $\text{sign}(\psi_l \psi_{l+2}) = -1$, induced by its occurrence in the energy functional (2): The energy is minimized by choosing $A_3 \psi_l \psi_{l+2} < 0$. Using the same argument, A_3 negative (Fig. 1, left panel) requires the signs of the ψ_l to be identical. The energy contribution of the pair-exchange term is generally negative, and its maximization forces ψ_l to be real.

The existence of sign changes has profound effects on the properties of superposed ground states. The first order coherence $g_1 = \frac{1}{2} \langle \hat{a}_0^\dagger \hat{a}_1 + \hat{a}_1^\dagger \hat{a}_0 \rangle$ vanishes for any superposition $a |\phi\rangle + b |\Phi\rangle$ in the presence of oscillations in the even and odd sectors of ψ_l , i.e. for $A_3 > 0$, leading to well-defined fragmented ground states, in the sense that their degree of fragmentation $\mathfrak{F}$ in (5) is independent of the concrete realisation via the weights a and b . To the best of our knowledge, this mechanism of fragmentation

for positive pair-exchange coupling in a trapped scalar Bose gas has not been discussed before in the literature. For A_3 negative, on the other hand, the possibility of superpositions allowed by the ground state degeneracy causes the system to be arbitrarily tunable between fully fragmented and coherent states; this can only be avoided by lifting the degeneracy by a small but finite Ω .

We define the degree of fragmentation to be $\mathfrak{F} = 1 - |\lambda_0 - \lambda_1|/N$, where $\lambda_{0,1}$ are the eigenvalues of the single-particle density matrix $\rho_{\mu\nu}^{(1)} = \langle a_\mu^\dagger a_\nu \rangle$, given by $\lambda_{0,1}/N = \frac{1}{2} \pm \sqrt{\frac{1}{4} - \frac{1}{N^2}(N_0 N_1 - |\langle \hat{a}_0^\dagger \hat{a}_1 \rangle|^2)}$. They collapse into a single nonvanishing one, $\lambda = N$, as soon as the state becomes coherent, $\langle \hat{a}_0^\dagger \hat{a}_1 \rangle = \sqrt{N_0 N_1}$, and therefore $\mathfrak{F} \equiv 0$ for coherent states. The degree of fragmentation evaluates to

$$\mathfrak{F} = 1 - \frac{2}{N} \sqrt{|\langle \hat{a}_0^\dagger \hat{a}_1 \rangle|^2 + \left(\frac{N}{2} - \sum_{l=1}^N |\psi_l|^2 l \right)^2}, \quad (5)$$

where $\langle \hat{a}_0^\dagger \hat{a}_1 \rangle = \sum_{l=1}^N \psi_{l-1}^* \psi_l \sqrt{l(N-l+1)}$.

The degeneracy of the two many-body states is lifted by Ω , which connects ψ_l and $\psi_{l\pm 1}$, and resolves this degeneracy already for $\Omega = \mathcal{O}(1/N)$ [so that the energy contribution of the Ω term is still suppressed to $\mathcal{O}(1/N^2)$ compared to the A_3 term]. For $A_3 < 0$, turning on Ω then establishes a coherent state characterized by number fluctuations $\Delta N_0^2 \propto N$ (with a proportionality factor of order unity), while for $A_3 > 0$, fragmentation persists for $\Omega > 0$. We illustrate the change of $\mathfrak{F}$ with A_3 , for finite Ω , in Fig. 2. We have checked the robustness of the fragmentation obtained on the $A_3 > 0$ side upon increasing the interlevel coupling up to $\Omega \sim \mathcal{O}(N^0)$, as well as for variations of the values of the single-particle energies ϵ_0 and ϵ_1 to the same order. The degree of fragmentation is therefore stable for small perturbations on the single-particle level, different to what was found in [16, 17], where the origin of fragmentation is distinct from our interaction-couplings based mechanism. Finally, because the energy contribution of the A_3 term is negative, for a concrete realization with sufficiently small A_3 the fragmented state has to be at a (local) minimum of the energy in the parameter space of the orbitals Ψ_i . The latter may be determined by a variational ansatz, e.g., by using the ellipsoid half-axes of harmonic oscillator trial wave functions as the orbital parameters [20, 21].

We now consider the continuum limit of slowly varying ψ_l [4], which allows for an analytical solution in the $\Omega \rightarrow 0$ limit. Employing the definition $j \equiv l - \frac{N}{2}$, and expanding c_j, d_j to quadratic order in j/N , neglecting $\mathcal{O}(1/N)$ terms, we obtain a differential equation for $|\psi(j)|$,

$$\left[-\frac{|A_3|N^2}{2} \frac{\partial^2}{\partial j^2} + \frac{|A_3|}{2R(\{A_i\})} (j - \mathfrak{S})^2 \right] |\psi| = E' |\psi|. \quad (6)$$




FIG. 2: The degree of fragmentation $\mathfrak{F}$ in Eq. (5) across the transition from coherent to fragmented states at $A_3 = 0$, for small but finite $\Omega = 1/N$. The blue dots represent the fragmentation from the numerical solution of (3); the solid red line is the continuum approximation from (9) in the $\Omega \rightarrow 0$ limit [the agreement, though not visible on the scale of the figure, has also been verified for the $A_3 < 0$ side]. The inset shows the first-order coherence, with the continuum result $g_1 = \sqrt{N^2/4 - \mathfrak{S}^2} \exp[-1/(4N\sqrt{R})]$.

Here, E' differs from E by a constant, and the location of the distribution center, the *shift*, is given by [24]

$$\mathfrak{S} = \frac{N(A_1 - A_2)/2 + \epsilon_0 - \epsilon_1}{A_1 + A_2 + 2|A_3| - A_4}. \quad (7)$$

The solution of lowest energy of the equation for $|\psi(j)|$ in (6) is a Gaussian centered at $\mathfrak{S}$, cf. Fig. 1, and of width $\sigma_{\text{osc}} = \sqrt{NR}^{1/4}(\{A_i\})$, provided that

$$R(\{A_i\}) = \frac{|A_3|}{A_1 + A_2 + 2|A_3| - A_4} \quad (8)$$

is positive. Note that particle number [to $\mathcal{O}(1/N)$] and single-particle energies do not enter R , in distinction to the corresponding criterion in the double-well case [4, 14]. For either fragmented ($A_3 > 0$) or coherent states ($A_3 < 0$), a necessary condition besides R being positive, is that the shift (7) fulfills $|\mathfrak{S}| \ll \frac{N}{2}$; in particular, a maximally fragmented state has $\mathfrak{S} = 0$. In the continuum limit, using the Gaussian shape of the $\psi(j)$ distribution, it may be shown that the degree of fragmentation is given by

$$\mathfrak{F} = \begin{cases} 1 - \frac{|\mathfrak{S}|}{N/2} & A_3 \geq 0, \\ 1 - e^{-\frac{1}{4N\sqrt{R}}} \sqrt{1 - \left(\frac{\mathfrak{S}}{N/2}\right)^2 \left(1 - e^{-\frac{1}{2N\sqrt{R}}}\right)} & A_3 < 0. \end{cases} \quad (9)$$

The above analytical approximation accurately reproduces the numerically obtained result, cf. Fig. (2). Note, provided we have the scaling $\epsilon_1 - \epsilon_0 = CN^\gamma$ for large N , with $\gamma \leq 1$ and C constants, assuming that all A_i scale identically and $A_1 \neq A_2$ when $C \neq 0$, we conclude from Eq. (7) that fragmentation persists for $N \rightarrow \infty$. This is a remarkable feature of the present two-mode model.

In case that R becomes negative, or diverges, in the nondegenerate case $c_0 \neq c_N$ we simply obtain Fock states

corresponding to the lower two-mode state, i.e. either $|N, 0\rangle$ or $|0, N\rangle$. The degenerate case leads to coherent superpositions of macroscopically distinct quantum states, for which the original single peak in the ψ_l distribution splits into two located around $l = 0$ and $l = N$. The crossover is illustrated in Fig. 3, by varying A_4 in (8), approaching the divergence of R . It appears that a fine-tuning of parameters is necessary, illustrating the sensitivity of the superposed states to parameter fluctuations [15].




FIG. 3: Crossover to a coherent superposition of degenerate macroscopically distinct quantum states for diverging R [Eq. (8)], using a maximally fragmented state with $\mathfrak{S} = 0$, upon variation of A_4 across the pole of R . Dark blue to light green (dark to light gray) indicates increasing A_4 ; $\Omega = 0$.

In conclusion, we have shown that adjusting the four interaction couplings in a two-mode model by employing the inhomogeneity of the single-particle orbitals in a trap, in conjunction with a generally anisotropic and nonlocal interaction, leads to fragmented ground states whose degree of fragmentation is not sensitive to perturbations on the single-particle level. The physical origin of this interaction-based fragmentation is the pair-exchange process of bosons in a trap, not occurring in the continuum. Pair exchanges cause oscillations in the distribution of the many-body wavefunction amplitudes, which as a consequence entail vanishing first-order phase coherence for positive pair-exchange coupling.

This research work was supported by the DFG under grant No. FI 690/3-1.

[1] O. Penrose and L. Onsager, Phys. Rev. **104**, 576 (1956).
[2] C. J. Pethick and L. P. Pitaevskiĭ, Phys. Rev. A **62**, 033609 (2000).
[3] P. Nozières and D. Saint James, J. Physique **43**, 1133 (1982).
[4] R. W. Spekkens and J. E. Sipe, Phys. Rev. A **59**, 3868 (1999).
[5] D. Jaksch *et al.*, Phys. Rev. Lett. **81**, 3108 (1998); M. Greiner *et al.*, Nature **415**, 39 (2002).
[6] For a summarizing review, see E. J. Mueller, T.-L. Ho, M. Ueda, and G. Baym, Phys. Rev. A **74**, 033612 (2006).

[7] T.-L. Ho and S. K. Yip, Phys. Rev. Lett. **84**, 4031 (2000).
[8] J. Dukelsky and P. Schuck, Phys. Rev. Lett. **86**, 4207 (2001).
[9] O. E. Alon and L. S. Cederbaum, Phys. Rev. Lett. **95**, 140402 (2005); L. S. Cederbaum, A. I. Streltsov, and O. E. Alon, Phys. Rev. Lett. **100**, 040402 (2008).
[10] N. K. Wilkin, J. M. F. Gunn, and R. A. Smith, Phys. Rev. Lett. **80**, 2265 (1998).
[11] See also the discussion in [2] of the example for fragmentation given in [10].
[12] G. J. Milburn, J. Corney, E. M. Wright, and D. F. Walls, Phys. Rev. A **55**, 4318 (1997).
[13] Such superposed macroscopic quantum states are sometimes referred to as “Schrödinger cats”, e.g. in [14, 15]. For clarity, we retain the explicit description of these states.
[14] T.-L. Ho and C. V. Ciobanu, J. Low Temp. Phys. **135**, 257 (2004).
[15] J. I. Cirac, M. Lewenstein, K. Mølmer, and P. Zoller, Phys. Rev. A **57**, 1208 (1998).
[16] R. Kanamoto, H. Saito, and M. Ueda, Phys. Rev. Lett. **94**, 090404 (2005).
[17] A. D. Jackson, G. M. Kavoulakis, and M. Magiropoulos, Phys. Rev. A **78**, 063623 (2008).
[18] A. J. Leggett, Rev. Mod. Phys. **73**, 307 (2001).
[19] A possible realization of finite Ω is to use two hyperfine states coupled by a two-photon Raman transition. Counterpropagating laser beams with effectively space-dependent Rabi frequency $\Omega_{\text{Rabi}}(\mathbf{r})$ lead to different spatial orbitals for the effective internal states [C. Zimmermann, private communication].
[20] For an illustrative example, we use as the two states the ground and an excited state of a harmonic oscillator along z , $\Psi_0(\rho, z) \equiv (a\sqrt{b}\pi^{\frac{3}{4}})^{-1} \exp[-\rho^2/2a^2 - z^2/2b^2]$ and $\Psi_1(\rho, z) \equiv (ab\sqrt{b/2}\pi^{\frac{3}{4}})^{-1} z \exp[-\rho^2/2a^2 - z^2/2b^2]$ (assuming cylindrical symmetry, where a and b are the half-axes of an ellipsoidal cloud [21]). Writing the dipole-dipole interaction as $V(\mathbf{r}) = \frac{3g_d}{4\pi|\mathbf{r}|^3}(1 - 3z^2/|\mathbf{r}|^2)$, where g_d is the dipole coupling constant [22], and adding the usual contact term, we obtain $A_1 = \frac{\sigma}{(\sqrt{2\pi}a)^3}[g + g_d f_1(\sigma)]$, $A_2 = \frac{3}{4} \frac{\sigma}{(\sqrt{2\pi}a)^3}[g + g_d f_2(\sigma)]$, $A_3 = \frac{1}{2} \frac{\sigma}{(\sqrt{2\pi}a)^3}[g + g_d f_3(\sigma)]$, and $A_4 = 2A_1$, where the $f_i(\sigma)$ are functions depending on the aspect ratio $\sigma = a/b$. They allow to sample the anisotropy and long-range character of the dipole-dipole interaction over the whole range from oblate ($\sigma > 1$) to prolate ($\sigma < 1$), while with contact interactions only, the ratios of the A_i are all constant [23]. Explicitly, $f_1 = \frac{1}{\sigma^2 - 1}[2\sigma^2 + 1 - 3\sigma^2 \text{ath}(\sigma)]$, $f_2 = \frac{1}{(\sigma^2 - 1)^3}[2\sigma^6 + 4\sigma^4 + 8\sigma^2 + 1 - \sigma^2(7 + 4\sigma^2 + 4\sigma^4)\text{ath}(\sigma)]$ and $f_3 = \frac{1}{(\sigma^2 - 1)^2}[2\sigma^4 - 10\sigma^2 - 1 + 9\sigma^2 \text{ath}(\sigma)]$, where $\text{ath}(\sigma) \equiv \arctan[\sqrt{\sigma^2 - 1}]/\sqrt{\sigma^2 - 1}$.
[21] Ø. Elgarøy and C. J. Pethick, Phys. Rev. A **59**, 1711 (1999).
[22] U. R. Fischer, Phys. Rev. A **73**, 031602(R) (2006).
[23] Going beyond the dominance of contact interactions in monatomic gases, by tuning the relative contributions of contact and dipole-dipole interactions using Feshbach resonances, has been experimentally demonstrated in T. Lahaye *et al.*, Nature **448**, 672 (2007).
[24] We observe here that the authors of [21] were in fact considering what we call shifted coherent states, which have only one eigenvalue of the single-particle density matrix, instead of fragmented states.